

### Contents lists available at ScienceDirect

## Asian Pacific Journal of Tropical Disease

journal homepage: www.elsevier.com/locate/apjtd



Document heading

doi: 10.1016/S2222-1808(14)60701-5

© 2014 by the Asian Pacific Journal of Tropical Disease. All rights reserved.

# Cysticercosis in returning traveler, a rare problem but really existed

Sora Yasri<sup>1\*</sup>, Viroj Wiwantikit<sup>2,3,4,5</sup>

<sup>1</sup>KMT Primary Care Center, Bangkok Thailand

<sup>2</sup>Hainan Medical University, China

<sup>3</sup>Joseph Ayobabalola University, Nigeria

<sup>4</sup>Dr DY Patil Medical University, India

<sup>5</sup>Faculty of Medicine, University of Nis, Serbia

To the editor,

Sir, the tropical infection among returning traveler is the present focus in tropical medicine and traveler medicine. The problem is usually due to the improved transportation and globalization at present. There are many reports on the tropical infections in returning travelers. The common reported infections include dengue, malaria etc. Of interest, a rare reported infection but really existed is the cysticercosis. Indeed, cysticercosis is a tapeworm infection and it requires long term for tissue infiltration and longer term to be detected. Here, the authors tried searching on the international records on this problem, using PubMed Search, and can determine 9 reports on this topic[1-8]. In those 9 reports, only 9 patients returning from endemic areas, South (especially for India) and Southeast Asia are reported[1,2,4]. For all cases, the accidental findings of parasite are noted. The unexplained mass lesion is the main chief complaints. Of interest, there are only a few reports on this issue. There are many questions on the situation. First, due to the long natural history of infection, the exact site that the patient got the infection is usually hard to identify. Sometimes, the patients might travel to more than one country. In addition, it is not possible to clear cut the episode of having get infection[9]. The history of food intake is only the presumptive diagnosis. Second, since the cysticercosis can be silent infection, sometimes, it can be misdiagnosed as neoplasia[1]. Hence, it is no doubt for the under reporting situation. Nevertheless, it can hereby conclude that cysticercosis in returning traveler is an existed problem. As we see, many new reports emerge within the few years implying the worldwide consideration on this existed problem. Finally, it should be repeatedly mentioned that cysticercosis in returning traveler is a rare

problem but really existed.

### **Conflict of interest statement**

We declare that we have no conflict of interest.

### References

- Salzer HJF, Hoenigl M, Valentin T, Zollner-Schwetz I, Krause R. Neurocysticercosis in a European traveler returning from India primarily suspected as neoplasia. *J Travel Med* 2013; 20(5): 333– 335
- [2] Kobayashi K, Nakamura-Uchiyama F, Nishiguchi T, Isoda K, Kokubo Y, Ando K, et al. Rare case of disseminated cysticercosis and taeniasis in a Japanese traveler after returning from India. Am J Trop Med Hyg 2013; 89(1): 58-62.
- [3] Del Brutto OH. Neurocysticercosis in Western Europe: a reemerging disease? *Acta Neurol Belg* 2012; **112**(4): 335–343.
- [4] Leshem E, Kliers I, Bakon M, Gomori M, Karplus R, Schwartz E. Neurocysticercosis in travelers: a nation-wide study in Israel. J Travel Med 2011; 18(3): 191-197.
- [5] Doherty T, Wright S. Cysticercosis in a female Nicaraguan traveler. J Travel Med 2010; 17(4): 288–289.
- [6] Croft AM, Flores AA, López HZ. Cysticercosis in a female Nicaraguan traveler. J Travel Med 2007; 14(5): 349–351.
- [7] Schantz PM, Wilkins PP, Tsang VC. A case of neurocysticercosis in a traveler to Latin America. Am J Trop Med Hyg 1999; 61(1): 1.
- [8] Chatel G, Gulletta M, Scolari C, Bombana E, El-Hamad I, Matteelli A, et al. Neurocysticercosis in an Italian traveler to Latin America. Am J Trop Med Hyg 1999; 60(2): 255–256.
- [9] Del Brutto OH, Nash TE, Garcia HH. Cysticerci-related single parenchymal brain enhancing lesions in non-endemic countries. J Neurol Sci 2012; 319(1-2): 32-36.